

Items Available Only in NEW location at MEB M4450

| Name | Size | Storage |
|---|-----------------|---------|
| Bovine Pituitary Extract (BPE) | 25 mg | -20G |
| Platinum™ Hot Start PCR Master Mix (2X) | 200 reactions | -20G |
| Smooth Muscle Growth Supplement (SMGS) | 25 mL | -20G |
| M-MLV Reverse Transcriptase Buffer | 1 mL | -20G |
| DNA-free™ DNA Removal Kit | 50 reactions | -20G |
| Human Keratinocyte Growth Supplement (HKGS) | 5 mL | -20G |
| bFGF Recombinant Human Protein | 10 µg | -20G |
| AccuPrime™ SuperMix II | 1,000 reactions | -20G |
| SuperScript™ III Reverse Transcriptase | 2,000 units | -20G |
| Phusion High-Fidelity DNA Polymerase (2 U/µL) | 100 units | -20G |
| Phusion Hot Start II DNA Polymerase (2 U/µL) | 100 units | -20G |
| PageRuler™ Plus Prestained Protein Ladder, 10 to 250 kDa | 2 x 250 µL | -20G |
| SuperScript™ First-Strand Synthesis System for RT-PCR | 50 reactions | -20G |
| SuperScript™ II Reverse Transcriptase | 10,000 units | -20G |
| SuperScript™ III First-Strand Synthesis System | 50 reactions | -20G |
| SuperScript™ III Reverse Transcriptase | 10,000 units | -20G |
| SuperScript™ VILO™ cDNA Synthesis Kit | 50 reactions | -20G |
| T4 DNA Ligase (1 U/µL) | 100 units | -20G |
| T4 DNA Ligase (5 U/µL) | 250 units | -20G |
| ThermoScript™ RT-PCR System for First-Strand cDNA Synthesis | 100 reactions | -20G |
| TrackIt™ 1 Kb Plus DNA Ladder | 50 µg | -20G |
| TrackIt™ 100 bp DNA Ladder | 50 µg | -20G |
| 1 Kb Plus DNA Ladder | 250 µg | -20G |
| dNTP Mix (10 mM ea) | 1 mL | -20G |
| dNTP Mix (10 mM ea) | 100 µL | -20G |
| 100 bp DNA Ladder | 50 µg | -20G |
| dNTP Set (100 mM) | 4 x 250 µL | -20G |
| N-2 Supplement (100X) | 5 mL | -20G |
| DNase I | 20,000 units | -20G |
| DNase I, Amplification Grade | 100 units | -20G |
| Geneticin™ Selective Antibiotic (G418 Sulfate) (50 mg/mL) | 20 mL | -20G |
| M-MLV Reverse Transcriptase (200 U/µL) | 40,000 units | -20G |
| MagicMark™ XP Western Protein Standard | 250 µL | -20G |
| Novex™ Sharp Pre-stained Protein Standard | 2 x 250 µL | -20G |
| Oligo(dT)12-18 Primer | 25 µg | -20G |
| Oligo(dT)20 Primer | 15 µg | -20G |
| PCR SuperMix | 100 reactions | -20G |
| Platinum™ Pfx DNA Polymerase | 100 reactions | -20G |
| Platinum™ Taq DNA Polymerase | 600 reactions | -20G |
| Platinum™ Taq DNA Polymerase | 120 reactions | -20G |
| Platinum™ Taq DNA Polymerase High Fidelity | 100 reactions | -20G |
| Platinum™ PCR SuperMix High Fidelity | 100 reactions | -20G |
| ProLong™ Gold Antifade Mountant | 10 mL | -20G |
| Proteinase K (Fungal) | 100 mg | -20G |

| | | |
|--|----------------|------|
| B-27™ Supplement (50X), serum free | 10 mL | -20G |
| B-27™ Supplement (50X), minus vitamin A | 10 mL | -20G |
| Power SYBR™ Green PCR Master Mix | 1 x 5 mL | -20G |
| Taq DNA Polymerase, native | 100 units | -20G |
| Taq DNA Polymerase, native | 500 units | -20G |
| Taq DNA Polymerase, recombinant | 100 units | -20G |
| Taq DNA Polymerase, recombinant | 500 units | -20G |
| B-27™ Supplement, minus insulin | 10 mL | -20G |
| BenchMark™ Pre-stained Protein Ladder | 2 x 250 µL | -20G |
| BenchMark™ Protein Ladder | 2 x 250 µL | -20G |
| TaqMan™ Fast Advanced Master Mix | 1 x 5 mL | -20G |
| RNaseOUT™ Recombinant Ribonuclease Inhibitor | 5,000 units | -20G |
| Ribonuclease H | 30 units | -20G |
| StemPro™ Accutase™ Cell Dissociation Reagent | 100 mL | -20G |
| DTT, 1M | 1 mL | -20G |
| TURBO DNA-free™ Kit | 50 reactions | -20G |
| High-Capacity cDNA Reverse Transcription Kit | 200 reactions | -20G |
| AccuPrime™ Pfx SuperMix | 200 reactions | -20G |
| SlowFade™ Diamond Antifade Mountant | 5 x 2 mL | -20G |
| ProLong™ Diamond Antifade Mountant | 5 x 2 mL | -20G |
| SlowFade™ Diamond Antifade Mountant with DAPI | 5 x 2 mL | -20G |
| ProLong™ Diamond Antifade Mountant with DAPI | 5 x 2 mL | -20G |
| Puromycin Dihydrochloride | 10 x 1 mL | -20G |
| FastDigest DpnI | 100 reactions | -20G |
| B-27™ Supplement (50X), custom | 10 mL | -20G |
| SuperScript™ IV First-Strand Synthesis System | 50 reactions | -20G |
| AccuPrime™ SuperMix II | 200 reactions | -20G |
| Oligo(dT)18 Primer | 120 µL | -20G |
| Fluo-4 NW Calcium Assay Kit | 10 microplates | -20H |
| Trypsin-EDTA (0.05%), phenol red | 500 mL | -20H |
| Trypsin-EDTA (0.05%), phenol red | 100 mL | -20H |
| Trypsin-EDTA (0.25%), phenol red | 500 mL | -20H |
| Trypsin-EDTA (0.5%), no phenol red | 100 mL | -20H |
| Antibiotic-Antimycotic (100X) | 100 mL | -20H |
| Penicillin-Streptomycin-Glutamine (100X) | 100 mL | -20H |
| Fetal Bovine Serum, certified, US origin | 500 mL | -20H |
| Fetal Bovine Serum, qualified, US origin | 500 mL | -20H |
| Fetal Bovine Serum, qualified, USDA-approved regions | 500 mL | -20H |
| Fetal Bovine Serum, qualified, heat inactivated, US origin | 500 mL | -20H |
| Gibco™ Amphotericin B | 20 mL | -20H |
| Horse Serum, heat inactivated, New Zealand origin | 500 mL | -20H |
| KnockOut™ Serum Replacement | 500 mL | -20H |
| Newborn Calf Serum, New Zealand origin | 500 mL | -20H |
| Penicillin-Streptomycin (10,000 U/mL) | 100 mL | -20H |
| L-Glutamine (200 mM) | 100 mL | -20H |
| Trypsin-EDTA (0.25%), phenol red | 100 mL | -20H |
| iBlot™ 2 Transfer Stacks, PVDF, mini | 10 stacks | RTA |

| | | |
|--|------------|-----|
| iBind™ Flex Cards | 10 cards | RTA |
| Restore™ Western Blot Stripping Buffer | 500 mL | RTA |
| Pierce™ BCA Protein Assay Kit | 1 L | RTA |
| iBlot™ 2 Transfer Stacks, PVDF, regular size | 10 stacks | RTA |
| PureLink™ HiPure Plasmid Filter Maxiprep Kit | 10 preps | RTA |
| PureLink™ HiPure Plasmid Filter Midiprep Kit | 25 preps | RTA |
| PureLink™ HiPure Plasmid Maxiprep Kit | 25 preps | RTA |
| PureLink™ HiPure Plasmid Maxiprep Kit | 10 preps | RTA |
| PureLink™ PCR Purification Kit | 50 preps | RTA |
| PureLink™ Quick Gel Extraction Kit | 50 preps | RTA |
| PureLink™ Quick Plasmid Miniprep Kit | 250 preps | RTA |
| PureLink™ Quick Plasmid Miniprep Kit | 50 preps | RTA |
| PureLink™ RNA Mini Kit | 50 preps | RTA |
| Restore™ PLUS Western Blot Stripping Buffer | 500 mL | RTA |
| SuperSignal™ West Pico PLUS Chemiluminescent Substrate | 200 mL | RTA |
| UltraPure™ DEPC-Treated Water | 4 x 100 mL | RTB |
| UltraPure™ DNase/RNase-Free Distilled Water | 500 mL | RTB |
| PBS (10X), pH 7.4 | 500 mL | RTB |
| TrypLE™ Express Enzyme (1X), phenol red | 500 mL | RTB |
| TrypLE™ Express Enzyme (1X), phenol red | 100 mL | RTB |
| DPBS, no calcium, no magnesium | 500 mL | RTB |
| DPBS, no calcium, no magnesium | 1,000 mL | RTB |
| Distilled Water | 1,000 mL | RTB |
| HBSS, calcium, magnesium, no phenol red | 1,000 mL | RTB |
| HBSS, calcium, magnesium, no phenol red | 500 mL | RTB |
| HBSS, no calcium, no magnesium | 500 mL | RTB |
| HBSS, no calcium, no magnesium, no phenol red | 500 mL | RTB |
| PBS, pH 7.4 | 500 mL | RTB |
| ACK Lysing Buffer | 100 mL | RTB |
| TrypLE™ Express Enzyme (1X), no phenol red | 500 mL | RTB |
| SYBR™ Safe DNA Gel Stain | 400 µL | RTB |
| TrypLE™ Express Enzyme (1X), no phenol red | 100 mL | RTB |
| Trypan Blue Solution, 0.4% | 100 mL | RTC |
| UltraPure™ 0.5M EDTA, pH 8.0 | 4 x 100 mL | RTC |
| UltraPure™ Agarose | 500 g | RTC |
| UltraPure™ Agarose | 100 g | RTC |
| UltraPure™ TAE Buffer, 10X | 1 L | RTC |
| XCell SureLock™ Mini-Cell | 1 unit | RTC |
| XCell II™ Blot Module | 1 unit | RTC |
| Gel Loading Tips, standard, round | 200/pack | RTC |
| Geneticin™ Selective Antibiotic (G418 Sulfate), Powder | 5 g | RTC |
| Gentamicin (50 mg/mL) | 10 mL | RTC |
| GlutaMAX™ Supplement | 100 mL | RTC |
| Invitrolon™ PVDF/Filter Paper Sandwich, 0.45 µm pore size, 8.3 cm x 7.3 cm | 20/pack | RTC |
| LB Broth Base | 500 g | RTC |
| MicroAmp™ Fast 8-Tube Strip, 0.1 mL | 125 strips | RTC |
| MicroAmp™ Fast Optical 96-Well Reaction Plate, 0.1 mL | 10 plates | RTC |

| | | |
|--|------------|-----|
| MicroAmp™ Optical 8-Cap Strips | 300 strips | RTC |
| MicroAmp™ Optical Adhesive Film | 100 covers | RTC |
| Nitrocellulose Pre-Cut Blotting Membranes, 0.45 µm pore size | 20/pack | RTC |
| Bolt™ Transfer Buffer (20X) | 1 L | RTC |
| Bottle Weight | 1 each | RTC |
| Cell Dissociation Buffer, enzyme-free, Hanks' Balanced Salt Solution | 100 mL | RTC |
| Cell Dissociation Buffer, enzyme-free, PBS | 100 mL | RTC |
| Countess™ Cell Counting Chamber Slides | 50 slides | RTC |
| DEPC-Treated Water | 1 x 100 mL | RTC |
| RNAlater™ Stabilization Solution | 100 mL | RTC |
| S.O.C. Medium | 10 x 10 mL | RTC |
| Sodium Bicarbonate 7.5% solution | 100 mL | RTC |
| Qubit™ Assay Tubes | 500 tubes | RTC |
| PureLink™ PCR Purification Kit | 250 preps | RTA |

Items Available Only on the 6th Floor of the MFRC

| Name | Size | Storage |
|---|----------|---------|
| DMEM, high glucose, no glutamine | 500 mL | CRD |
| DMEM, high glucose, pyruvate | 500 mL | CRD |
| DMEM, low glucose, pyruvate | 500 mL | CRD |
| DMEM, no glucose | 500 mL | CRD |
| DMEM, high glucose | 500 mL | CRD |
| DMEM, high glucose | 1,000 mL | CRD |
| DMEM, high glucose, no glutamine, no phenol red | 500 mL | CRD |
| RPMI 1640 Medium | 500 mL | CRD |
| RPMI 1640 Medium, HEPES | 500 mL | CRD |
| RPMI 1640 Medium, no phenol red | 500 mL | CRD |
| RPMI 1640 Medium (ATCC modification) | 500 mL | CRD |
| RPMI 1640 Medium | 1,000 mL | CRD |
| KnockOut™ DMEM | 500 mL | CRE |
| Neurobasal™ Medium | 500 mL | CRE |
| DMEM, no glucose, no glutamine, no phenol red | 500 mL | CRE |
| DMEM/F-12, HEPES | 500 mL | CRE |
| DPBS, calcium, magnesium | 500 mL | CRE |
| Ham's F-12 Nutrient Mix | 500 mL | CRE |
| Ham's F-12K (Kaighn's) Medium | 500 mL | CRE |
| IMDM | 500 mL | CRE |
| MEM | 500 mL | CRE |
| MEM α, no nucleosides | 500 mL | CRE |
| MEM, NEAA, no glutamine | 500 mL | CRE |
| McCoy's 5A (Modified) Medium | 500 mL | CRE |
| Medium 199, HEPES | 500 mL | CRE |
| Opti-MEM™ I Reduced Serum Medium | 500 mL | CRE |
| Opti-MEM™ I Reduced Serum Medium | 100 mL | CRE |
| RPMI 1640 Medium, no glutamine | 500 mL | CRE |
| DMEM/F-12 | 500 mL | CRE |
| William's E Medium, no glutamine | 500 mL | CRE |

| | | |
|---|-----------------|-----|
| PowerUp™ SYBR™ Green Master Mix | 1 mL | CRF |
| iBind™ Flex Solution Kit | 1 kit | CRF |
| TaqMan™ Universal Master Mix II, with UNG | 1 x 5 mL | CRF |
| TaqMan™ Gene Expression Master Mix | 1 x 1 mL | CRF |
| NuPAGE™ Sample Reducing Agent (10X) | 10 mL | CRF |
| GeneScan™ 600 LIZ™ dye Size Standard v2.0 | 800 reactions | CRF |
| Lipofectamine™ 3000 Transfection Reagent | 1.5 mL | CRF |
| Novex™ 4-20% Tris-Glycine Mini Gels, WedgeWell™ format, 15-well | 10 gels (1 box) | CRF |
| SuperSignal™ West Femto Maximum Sensitivity Substrate | 100 mL | CRF |
| Pierce™ ECL Plus Western Blotting Substrate | 100 mL | CRF |
| Halt™ Protease Inhibitor Cocktail (100X) | 24 x 100 µL | CRF |
| TRizol™ Reagent | 100 mL | CRF |
| TRizol™ Reagent | 200 mL | CRF |
| TaqMan™ Genotyping Master Mix | 1 x 10 mL | CRF |
| TaqMan™ Genotyping Master Mix | 1 x 1 mL | CRF |
| TaqMan™ Universal Master Mix II, with UNG | 1 x 1 mL | CRF |
| Versene Solution | 100 mL | CRF |
| Sodium Pyruvate (100 mM) | 100 mL | CRF |
| Insulin-Transferrin-Selenium (ITS -G) (100X) | 10 mL | CRF |
| MEM Non-Essential Amino Acids Solution (100X) | 100 mL | CRF |
| BlueJuice™ Gel Loading Buffer (10X) | 3 x 1 mL | CRF |
| HEPES (1 M) | 100 mL | CRF |
| Hygromycin B (50 mg/mL) | 20 mL | CRF |
| Lipofectamine™ 2000 Transfection Reagent | 1.5 mL | CRF |
| Lipofectamine™ 2000 Transfection Reagent | 0.75 mL | CRF |
| Lipofectamine™ LTX Reagent with PLUS™ Reagent | 1 mL | CRF |
| Lipofectamine™ RNAiMAX Transfection Reagent | 0.75 mL | CRF |
| NuPAGE™ Antioxidant | 15 ml | CRF |
| NuPAGE™ LDS Sample Buffer (4X) | 10 mL | CRF |
| NuPAGE™ MES SDS Running Buffer (20X) | 500 mL | CRF |
| NuPAGE™ MOPS SDS Running Buffer (20X) | 500 mL | CRF |
| NuPAGE™ 10% Bis-Tris Protein Gels, 1.0 mm, 10-well | 10 gels (1 box) | CRF |
| NuPAGE™ 10% Bis-Tris Protein Gels, 1.0 mm, 15-well | 10 gels (1 box) | CRF |
| NuPAGE™ 4-12% Bis-Tris Protein Gels, 1.5 mm, 10-well | 10 gels (1 box) | CRF |
| NuPAGE™ 4-12% Bis-Tris Protein Gels, 1.5 mm, 15-well | 10 gels (1 box) | CRF |
| NuPAGE™ Sample Reducing Agent (10X) | 250 µL | CRF |
| NuPAGE™ Transfer Buffer (20X) | 1 L | CRF |
| Gibco™ 2-Mercaptoethanol | 50 mL | CRF |
| TaqMan™ Gene Expression Master Mix | 1 x 5 mL | CRF |
| SeeBlue™ Plus2 Pre-stained Protein Standard | 500 µL | CRF |
| Lipofectamine™ 3000 Transfection Reagent | 0.75 mL | CRF |
| RIPA Lysis and Extraction Buffer | 100 mL | CRF |
| Pierce™ Quantitative Fluorometric Peptide Assay | 500 assays | CRF |
| Bovine Albumin Fraction V (7.5% solution) | 100 mL | CRF |
| MagniSort™ Mouse CD4 Positive Selection Kit | 200 tests | CRF |
| MagniSort™ Mouse T cell Enrichment Kit | 200 tests | CRF |
| eBioscience™ Intracellular Fixation & Permeabilization Buffer Set | 1 kit | CRF |

| | | |
|--|--------------|--------|
| TaqMan™ Universal Master Mix II, no UNG | 1 x 50 mL | CRF |
| IL-1 beta Mouse Uncoated ELISA Kit with Plates | 2 x 96 tests | CRF |
| Nunc™ EasYDish™, 100mm / 17mm | Case of 150 | MEB4TH |
| Nunc™ EasYDish™, 60mm | Case of 280 | MEB4TH |
| Nunc™ Cell-Culture Treated Multidishes, 24 well | Case of 75 | MEB4TH |
| Nunc™ Conical Sterile Polypropylene Centrifuge Tubes, 15mL, racked | Case of 500 | MEB4TH |
| Nunc™ Conical Sterile Polypropylene Centrifuge Tubes, 50mL, racked, 50mL | Case of 300 | MEB4TH |
| Nunc™ Serological Pipets, 5mL, paper peel | Case of 200 | MEB4TH |
| Nunc™ Serological Pipets, 10mL, paper peel | Case of 200 | MEB4TH |
| Nunc™ Serological Pipets, 25mL, paper peel | Case of 200 | MEB4TH |
| Nunc™ EasYFlask™ Cell Culture Flasks, T75, filter | Case of 100 | MEB4TH |
| Nunc™ EasYFlask™ Cell Culture Flasks, T175, solid cap | Case of 30 | MEB4TH |
| Nunc™ Cell-Culture Treated Multidishes, 6 well | Case of 75 | MEB4TH |
| Nunc™ Cell-Culture Treated Multidishes, 12 well | Case of 75 | MEB4TH |
| Essential 8™ Medium | 500 mL | |