ADAMS COUNTY

The information presented here represents data specific to Adams County, Wisconsin from 2001-2006, and is meant to complement the report, *The Burden of Suicide in Wisconsin*. Please refer to this report for technical notes and suicide prevention strategies. For counties with fewer than 20 suicides each year, data are aggregated for 2001-2006 and findings are presented in bullets rather than tables.

KEY FINDINGS

- ◆The suicide rate in Adams County for all years was higher than the state rate for all years (16.9 versus 11.8).
- ♦ Suicide represented a total of 547 years of potential life lost, at a rate of 476.8 for 2001-2006. This was more than the state rate (394.1 YPLL) for the same time period.
- ◆The greatest number of suicides occurred in individuals aged 35-54, which is the same at the state level. The greatest number of hospitalizations and emergency department visits occurred in those aged 35-44, which is older than at the state level.
- ◆The cost of inpatient hospitalizations and emergency department visits due to self-inflicted injury was over \$76,000 in 2006 alone.
- ♦ Firearms were the most frequently used method in Adams County suicides (48%) followed by poisoning (19%) and hanging, strangulation, or suffocation (19%).

Numbers and rates (per 100,000 population) of suicides, inpatient hospitalizations, and emergency department visits due to self-inflicted injuries, by age, Adams County, 2001-2006.

	Deaths		Inpatient Hospitalizations		Emergency Department Visit*	
Age Group	Number	Rate‡	Number	Rate‡	Number	Rate‡
5-14	I	8.0‡	5	40.1‡	5	48.7‡
15-24	Ι	8.1‡	14	112.9‡	21	201.6
25-34	0		14	102.9‡	21	184.2
35-44	5	27.4‡	24	131.5	23	152.4
45-54	5	25.7‡	9	46.3‡	10	60.7‡
55-64	2	10.8‡	10	53.9‡	I	6.4‡
65-74	3	20.1‡	2	13.4‡	2	16.0‡
75-84	3	39.5‡	I	13.2‡	0	
85+	0		0		0	
Total	20	16.9	79	66.1	83	82.9

^{*}Emergency department data are not available for 2001.