CHIPPEWA COUNTY

The information presented here represents data specific to Chippewa County, Wisconsin from 2001-2006, and is meant to complement the report, *The Burden of Suicide in Wisconsin*. Please refer to this report for technical notes and suicide prevention strategies. For counties with fewer than 20 suicides each year, data are aggregated for 2001-2006 and findings are presented in bullets rather than tables.

KEY FINDINGS

- ◆The suicide rate in Chippewa County for all years was higher than the state rate for all years (15.9 versus 11.8).
- ♦ Suicide represented a total of 1,541 years per life lost, at a rate of 472.1 for 2001-2006. This was more than the state rate (394.1 YPLL) for the same time period.
- ◆The greatest number of suicides occurred in individuals aged 45-54, while the greatest number of hospitalizations and emergency department visits occurred in younger individuals, aged 15-24. This finding is the same at the state level.
- ◆The cost of inpatient hospitalizations and emergency department visits due to self-inflicted injury was over \$870,000 in 2006 alone.
- ♦ Firearms were the most frequently used method in Chippewa County suicides (57%) followed by poisoning (23%) and hanging, strangulation, or suffocation (14%).

Numbers and rates (per 100,000 population) of suicides, inpatient hospitalizations, and emergency department visits due to self-inflicted injuries, by age, Chippewa County, 2001-2006.

	Deaths		Inpatient Hospitalizations		Emergency Department Visit*	
Age Group	Number	Rate‡	Number	Rate‡	Number	Rate‡
5-14	0		23	47.8	5	12.5‡
15-24	5	10.5‡	123	257.1	39	96.5
25-34	10	24.5‡	73	179.4	23	67.2
35-44	9	16.5‡	65	119.4	16	35.3‡
45-54	16	29.8‡	47	87.7	4	7.5‡
55-64	3	8.4‡	12	33.7‡	I	2.8‡
65-74	9	36.6‡	5	20.2‡	0	
75-84	2	I I.2‡	2	II.2‡	0	
85+	2	25.5‡	0		0	
Total	56	15.9	350	99.4	88	30.0

^{*}Emergency department data are not available for 2001.