GRANT COUNTY

The information presented here represents data specific to Grant County, Wisconsin from 2001-2006, and is meant to complement the report, *The Burden of Suicide in Wisconsin*. Please refer to this report for technical notes and suicide prevention strategies. For counties with fewer than 20 suicides each year, data are aggregated for 2001-2006 and findings are presented in bullets rather than tables.

KEY FINDINGS

- ◆The suicide rate in Grant County for all years was lower than the state rate for all years (5.6‡ versus 11.8).
- ♦ Suicide represented a total of 1,442 years of potential life lost, at a rate of 520.6 for 2001-2006. This was more than the state rate (394.1 YPLL) for the same time period.
- ◆The greatest number of suicides occurred in individuals aged 15-24, which is younger than the state as a whole. The greatest number of hospitalizations and emergency department visits occurred in the same age group, which is consistent with statewide findings.
- ◆The cost of inpatient hospitalizations and emergency department visits due to self-inflicted injury was over \$449,000 in 2006 alone.
- ♦ Firearms were the most frequently used method in Grant County suicides (53%) followed by hanging, strangulation, or suffocation (29%) and poisoning (18%).

Numbers and rates (per 100,000 population) of suicides, inpatient hospitalizations, and emergency department visits due to self-inflicted injuries, by age, Grant County, 2001-2006.

	Deaths		Inpatient Hospitalizations		Emergency Department Visit*	
Age Group	Number	Rate‡	Number	Rate‡	Number	Rate‡
5-14	0		7	19.0‡	П	36.3‡
15-24	7	11.5‡	73	119.9	55	108.0
25-34	3	10.0‡	42	139.8	18	72.I‡
35-44	4	10.3‡	34	87.3	10	31.3‡
45-54	I	2.4‡	23	54.6	5	14.1‡
55-64	I	3.3‡	13	43.5‡	4	15.8‡
65-74	I	4.4‡	7	30.8‡	2	10.5‡
75-84	0		0		0	
85+	0		I	13.4‡	I	15.9‡
Total	17	5.6 ‡	200	70.1	106	44.6

^{*}Emergency department data are not available for 2001.