GREEN LAKE COUNTY

The information presented here represents data specific to Green Lake County, Wisconsin from 2001-2006, and is meant to complement the report, *The Burden of Suicide in Wisconsin*. Please refer to this report for technical notes and suicide prevention strategies. For counties with fewer than 20 suicides each year, data are aggregated for 2001-2006 and findings are presented in bullets rather than tables.

KEY FINDINGS

- ◆The suicide rate in Green Lake County for all years was higher than the state rate for all years (14.7‡ versus 11.8).
- ♦ Suicide represented a total of 390 years of potential life lost, at a rate of 373.7 for 2001-2006. This was less than the state rate (394.1 YPLL) for the same time period.
- ♦The greatest number of suicides occurred in individuals aged 45-54, while the greatest number of hospitalizations and emergency department visits occurred in younger individuals, aged 15-24. This finding is the same at the state level.
- ♦The cost of inpatient hospitalizations and emergency department visits due to self-inflicted injury was over \$72,000 in 2006 alone.
- ♦ Firearms were the most frequently used method in Green Lake County suicides (53%) followed by hanging, strangulation, or suffocation (29%).

Numbers and rates (per 100,000 population) of suicides, inpatient hospitalizations, and emergency department visits due to self-inflicted injuries, by age, Green Lake County, 2001-2006.

	Deaths		Inpatient Hospitalizations		Emergency Department Visit*	
Age Group	Number	Rate‡	Number	Rate‡	Number	Rate‡
5-14	0		0		2	17.1‡
15-24	2	14.1‡	23	162.0	25	210.2
25-34	0		18	158.5‡	9	95.6‡
35-44	I	6.0‡	19	I I 4.0‡	11	80.4‡
45-54	6	33.1‡	7	38.6‡	6	39.2‡
55-64	4	29.8‡	2	14.9‡	0	
65-74	3	30.2‡	3	30.2‡	0	
75-84	I	12.8‡	3	38.5‡	I	15.4‡
85+	0		0		0	
Total	17	14.7‡	75	68.7	54	59.3

^{*}Emergency department data are not available for 2001.