LANGLADE COUNTY

The information presented here represents data specific to Langlade County, Wisconsin from 2001-2006, and is meant to complement the report, *The Burden of Suicide in Wisconsin*. Please refer to this report for technical notes and suicide prevention strategies. For counties with fewer than 20 suicides each year, data are aggregated for 2001-2006 and findings are presented in bullets rather than tables.

KEY FINDINGS

- ◆The suicide rate in Langlade County for all years was lower than the state rate for all years (7.1 versus 11.8).
- ◆Suicide represented a total of 470 years of potential life lost, at a rate of 411.2 for 2001-2006. This was more than the state rate (394.1 YPLL) for the same time period.
- ◆The greatest number of suicides occurred in individuals aged 35-44, while the greatest number of hospitalizations and emergency department visits occurred in younger individuals, aged 15-24. This finding is the same at the state level.
- ◆The cost of inpatient hospitalizations and emergency department visits due to self-inflicted injury was over \$128,000 in 2006 alone.
- ♦ Firearms were the most frequently used method in Langlade County suicides (78%) followed by hanging, strangulation, or suffocation (11%) and fire (11%).

Numbers and rates (per 100,000 population) of suicides, inpatient hospitalizations, and emergency department visits due to self-inflicted injuries, by age, Langlade County, 2001-2006.

	Deaths		Inpatient Hospitalizations		Emergency Department Visit*	
Age Group	Number	Rate‡	Number	Rate‡	Number	Rate‡
5-14	0		6	37.8‡	3	23.I‡
15-24	2	13.0‡	27	175.6	29	223.8
25-34	0		6	47.3‡	П	105.6‡
35-44	4	22.1‡	22	121.4	12	80.3‡
45-54	I	5.3‡	6	31.9‡	9	56.9‡
55-64	I	6.6‡	2	13.3‡	3	23.5‡
65-74	0		0		ı	10.2‡
75-84	I	I I.6‡	2	23.2‡	2	27.8‡
85+	0	-	0		0	
Total	9	7. 1‡	71	59.2	70	70.0

^{*}Emergency department data are not available for 2001.