## **OZAUKEE COUNTY**

The information presented here represents data specific to Ozaukee County, Wisconsin from 2001-2006, and is meant to complement the report, *The Burden of Suicide in Wisconsin*. Please refer to this report for technical notes and suicide prevention strategies. For counties with fewer than 20 suicides each year, data are aggregated for 2001-2006 and findings are presented in bullets rather than tables.

## **KEY FINDINGS**

- ◆The suicide rate in Ozaukee County for all years was lower than the state rate for all years (10.4 versus 11.8).
- ♦ Suicide represented a total of 1,660 years of potential life lost, at a rate of 346.7 for 2001-2006. This was less than the state rate (394.1 YPLL) for the same time period.
- ◆The greatest number of suicides occurred in individuals aged 35-54, while the greatest number of hospitalizations and emergency department visits occurred in younger individuals, aged 15-24. This finding is the same at the state level.
- ◆The cost of inpatient hospitalizations and emergency department visits due to self-inflicted injury was over \$715,000 in 2006 alone.
- ♦ Hanging, strangulation, or suffocation was the most frequently used method in Ozaukee County suicides (45%) followed by firearms (28%) and poisoning (21%).

Numbers and rates (per 100,000 population) of suicides, inpatient hospitalizations, and emergency department visits due to self-inflicted injuries, by age, Ozaukee County, 2001-2006.

	Deaths		Inpatient Hospitalizations		Emergency Department Visit*	
Age Group	Number	Rate‡	Number	Rate‡	Number	Rate‡
5-14	0		8	10.7‡	22	35.7
15-24	6	9.2‡	108	164.8	84	151.4
25-34	8	16.6‡	73	151.8	8	20.1‡
35-44	13	15.9‡	97	118.8	9	13.4‡
45-54	13	14.7‡	47	53.3	9	12.1‡
55-64	7	12.1‡	19	32.8‡	2	4.0‡
65-74	2	5.8‡	8	23.4‡	I	3.5‡
75-84	3	12.8‡	3	12.8‡	1	5.1‡
85+	I	I I.7‡	0		I	13.8‡
Total	53	10.4	363	75.3	137	33.9

<sup>\*</sup>Emergency department data are not available for 2001.