PRICE COUNTY

The information presented here represents data specific to Price County, Wisconsin from 2001-2006, and is meant to complement the report, *The Burden of Suicide in Wisconsin*. Please refer to this report for technical notes and suicide prevention strategies. For counties with fewer than 20 suicides each year, data are aggregated for 2001-2006 and findings are presented in bullets rather than tables.

KEY FINDINGS

◆The suicide rate in Price County for all years was higher than the state rate for all years (14.8⁺ versus 11.8).

- ◆Suicide represented a total of 405 years of potential life lost, at a rate of 474.7 for 2001-2006. This was more than the state rate (394.1 YPLL) for the same time period.
- ◆The greatest number of suicides occurred in individuals aged 45-54, while the greatest number of hospitalizations and emergency department visits occurred in younger individuals, aged 15-24. This finding is the same at the state level.
- ◆The cost of inpatient hospitalizations and emergency department visits due to self-inflicted injury was over \$20,000 in 2006 alone.
- ♦ Firearms were the most frequently used method in Price County suicides (71%) followed by hanging, strangulation, or suffocation (14%).

	Deaths		Inpatient Hospitalizations		Emergency Department Visit*	
Age Group	Number	Rate‡	Number	Rate‡	Number	Rate‡
5-14	0		4	34.7‡	2	21.3‡
15-24	I	8.9‡	25	223.3	15	I 58.9‡
25-34	2	23.1‡	8	92.3‡	8	112.3‡
35-44	2	14.6‡	П	80.4‡	5	44.4
45-54	6	38.2‡	12	76.3‡	2	15.1‡
55-64	2	16.9‡	5	42.2‡	0	
65-74	0		3	35.0‡	0	
75-84	I	15.9‡	0		I	19.1‡
85+	0		0		0	
Total	14	14.8 ‡	68	75.2	33	43.8

Numbers and rates (per 100,000 population) of suicides, inpatient hospitalizations, and emergency department visits due to self-inflicted injuries, by age, Price County, 2001-2006.

*Emergency department data are not available for 2001.

*Rates based on numbers of 20 or less are considered unstable due to random chance and should be interpreted with caution.