SHAWANO COUNTY

The information presented here represents data specific to Shawano County, Wisconsin from 2001-2006, and is meant to complement the report, *The Burden of Suicide in Wisconsin*. Please refer to this report for technical notes and suicide prevention strategies. For counties with fewer than 20 suicides each year, data are aggregated for 2001-2006 and findings are presented in bullets rather than tables.

KEY FINDINGS

◆The suicide rate in Shawano County for all years was lower than the state rate for all years (8.4 versus 11.8).

- ◆Suicide represented a total of 1,155 years of potential life lost, at a rate of 505.5 for 2001-2006. This was more than the state rate (394.1 YPLL) for the same time period.
- ◆The greatest number of suicides occurred in individuals aged 45-54, while the greatest number of hospitalizations and emergency department visits occurred in younger individuals, aged 15-24. This finding is the same at the state level.
- ♦The cost of inpatient hospitalizations and emergency department visits due to self-inflicted injury was over \$530,000 in 2006 alone.
- ◆Firearms were the most frequently used method in Shawano County suicides (71%) followed by poisoning (19%).

	Deaths		Inpatient Hospitalizations		Emergency Department Visit*	
Age Group	Number	Rate‡	Number	Rate‡	Number	Rate‡
5-14	0		17	50.4‡	17	61.0 †
15-24	4	12.7‡	78	247.7	76	286.8
25-34	3	10.9‡	33	119.7	37	161.9
35-44	3	8.0‡	41	109.5	28	90.1
45-54	6	17.1‡	23	65.6	12	40.6‡
55-64	0		12	42.4‡	3	12.5‡
65-74	2	9.7 ‡	4	19.7‡	2	۱۱. 9 ‡
75-84	3	20.1‡	I	6.7‡	0	
85+	0		2	32.3‡	0	
Total	21	8.4	211	89.8	175	89.1

Numbers and rates (per 100,000 population) of suicides, inpatient hospitalizations, and emergency department visits due to self-inflicted injuries, by age, Shawano County, 2001-2006.

*Emergency department data are not available for 2001.

‡Rates based on numbers of 20 or less are considered unstable due to random chance and should be interpreted with caution.