TAYLOR COUNTY

The information presented here represents data specific to Taylor County, Wisconsin from 2001-2006, and is meant to complement the report, *The Burden of Suicide in Wisconsin*. Please refer to this report for technical notes and suicide prevention strategies. For counties with fewer than 20 suicides each year, data are aggregated for 2001-2006 and findings are presented in bullets rather than tables.

KEY FINDINGS

◆The suicide rate in Taylor County for all years was higher than the state rate for all years (12.6⁺ versus 11.8).

- ◆Suicide represented a total of 327 years of potential life lost, at a rate of 299.3 for 2001-2006. This was less than the state rate (394.1 YPLL) for the same time period.
- ♦ The greatest number of suicides occurred in individuals aged 65-74, which was older than the state as a whole. The greatest number of hospitalizations and emergency department visits occurred in younger individuals, aged 15-24. This finding is the same at the state level.
- ♦The cost of inpatient hospitalizations and emergency department visits due to self-inflicted injury was over \$28,000 in 2006 alone.
- ◆Firearms were the most frequently used method in Taylor County suicides (67%) followed by poisoning (13%).

	Deaths		Inpatient Hospitalizations		Emergency Department Visit*	
Age Group	Number	Rate‡	Number	Rate‡	Number	Rate‡
5-14	0		5	30.3‡	2	14.8‡
15-24	0		35	213.4	27	196.5
25-34	2	15.2‡	13	99.1‡	3	27.5‡
35-44	2	11.0‡	9	49.4‡	5	33.4 †
45-54	2	11.2‡	4	22.5‡	4	26.5‡
55-64	2	17.4‡	2	17.4‡	0	
65-74	4	45.7‡	0		0	
75-84	3	46.9 ‡	I	15.6‡	0	
85+	0		0		0	
Total	15	I 2.6 ‡	69	61.8	41	44.0

Numbers and rates (per 100,000 population) of suicides, inpatient hospitalizations, and emergency department visits due to self-inflicted injuries, by age, Taylor County, 2001-2006.

*Emergency department data are not available for 2001.

‡Rates based on numbers of 20 or less are considered unstable due to random chance and should be interpreted with caution.