## **WASHINGTON COUNTY**

The information presented here represents data specific to Washington County, Wisconsin from 2001-2006, and is meant to complement the report, *The Burden of Suicide in Wisconsin*. Please refer to this report for technical notes and suicide prevention strategies. For counties with fewer than 20 suicides each year, data are aggregated for 2001-2006 and findings are presented in bullets rather than tables.

## **KEY FINDINGS**

- ◆The suicide rate in Washington County for all years was lower than the state rate for all years (9.2 versus 11.8).
- ♦ Suicide represented a total of 2,347 years of potential life lost, at a rate of 335.0 for 2001-2006. This was less than the state rate (394.1 YPLL) for the same time period.
- ◆The greatest number of suicides occurred in individuals aged 35-44, while the greatest number of hospitalizations and emergency department visits occurred in younger individuals, aged 15-24. This finding is the same at the state level.
- ◆The cost of inpatient hospitalizations and emergency department visits due to self-inflicted injury was over \$953,000 in 2006 alone.
- ♦ Firearms were the most frequently used method in Washington County suicides (47%) followed by hanging, strangulation, or suffocation (28%) and poisoning (22%).

Numbers and rates (per 100,000 population) of suicides, inpatient hospitalizations, and emergency department visits due to self-inflicted injuries, by age, Washington County, 2001-2006.

	Deaths		Inpatient Hospitalizations		Emergency Department Visit*	
Age Group	Number	Rate‡	Number	Rate‡	Number	Rate‡
5-14	0		19	17.9‡	17	19.3‡
15-24	12	13.0‡	133	144.3	60	77.2
25-34	7	7.9‡	52	58.5	16	21.7‡
35-44	19	14.5‡	74	56.3	31	28.3
45-54	17	14.6‡	62	53.4	12	I 2.2‡
55-64	8	10.7‡	13	I7.4‡	4	6.3‡
65-74	I	2.3‡	4	9.0 <del>‡</del>	0	
75-84	3	9.9‡	3	9.9 <del>‡</del>	0	
85+	I	8.4‡	0		0	
Total	68	9.2	360	51.7	140	24.0

<sup>\*</sup>Emergency department data are not available for 2001.