WAUPACA COUNTY

The information presented here represents data specific to Waupaca County, Wisconsin from 2001-2006, and is meant to complement the report, *The Burden of Suicide in Wisconsin*. Please refer to this report for technical notes and suicide prevention strategies. For counties with fewer than 20 suicides each year, data are aggregated for 2001-2006 and findings are presented in bullets rather than tables.

KEY FINDINGS

- ◆The suicide rate in Waupaca County for all years was lower than the state rate for all years (10.4 versus 11.8).
- ♦ Suicide represented a total of 1,128 years of potential life lost, at a rate of 390.9 for 2001-2006. This was less than the state rate (394.1 YPLL) for the same time period.
- ◆The greatest number of suicides occurred in individuals aged 35-54, while the greatest number of hospitalizations and emergency department visits occurred in younger individuals, aged 15-24. This finding is the same at the state level.
- ◆The cost of inpatient hospitalizations and emergency department visits due to self-inflicted injury was over \$376,000 in 2006 alone.
- ♦ Firearms were the most frequently used method in Waupaca County suicides (58%) followed by poisoning (24%) and hanging, strangulation, or suffocation (18%).

Numbers and rates (per 100,000 population) of suicides, inpatient hospitalizations, and emergency department visits due to self-inflicted injuries, by age, Waupaca County, 2001-2006.

	Deaths		Inpatient Hospitalizations		Emergency Department Visit*	
Age Group	Number	Rate‡	Number	Rate‡	Number	Rate‡
5-14	I	2.3‡	16	37.4‡	9	25.5‡
15-24	5	12.4‡	84	208.7	77	227.4
25-34	5	14.4‡	21	60.4‡	21	72.7
35-44	П	22.8‡	42	87.0	45	112.8
45-54	4	8.4‡	32	67.4	17	42.4‡
55-64	4	12.0‡	8	24.0‡	3	10.5‡
65-74	3	12.6‡	8	33.5‡	0	
75-84	0		1	5.1‡	I	6.1‡
85+	0		I	II.0‡	2	26.2‡
Total	33	10.4	213	71.1	175	69.9

^{*}Emergency department data are not available for 2001.