





Fatal Drug Combinations in OFR Cases

Out of 792 cases reviewed by Overdose Fatality Review (OFR) teams in Wisconsin since 2018, 681 had one or more drug types identified for the cause of death. Of these 681 cases, 488 overdose fatalities have been attributed to multiple substances (approximately 72 percent). Overall, fentanyl was the most common substance found in combination with other drugs. Out of the 681 cases reviewed, 148 cases involved a combination of fentanyl and cocaine, 107 cases involved a combination of fentanyl and heroin, and 103 cases involved a combination of fentanyl and methamphetamine.

Frequency of Fatal Drug Combinations in OFR Cases, 2018-2023


Each line represents a potential pairing of drugs that was involved in an overdose fatality. The thickness and color indicate the percentage of all OFR cases (where a substance was identified) in which that drug combination was present, as indicated on the death certificate. In fatalities in which more than two substances were involved, the substances were divided into unique combinations of two.



Single drug deaths are not displayed, but percentages are calculated out of ALL OFR cases where a substance could be identified.
N cases = 681. N multi-drug cases = 488.

Substances in All OFR Cases

Across all 792 cases, fentanyl was observed in the most overdose fatalities, recorded as present in more than half of the incidents. In both single and multiple drug deaths, fentanyl, cocaine, other opioids, and benzodiazepines were most likely to be involved. After benzodiazepines, methamphetamine were involved in the most deaths reviewed by OFR teams.



Methodology

Substances were grouped into categories based on terms contained in the cause of death or the description of the cause of death collected in the Wisconsin OFR database. Terms included both generic and brand names of substances. The analysis includes the most commonly listed substances listed among the causes of overdose deaths reviewed by OFR teams in Wisconsin. This visualization is not an exhaustive list of all drug combinations in OFR cases.

Once each case was classified into one or more categories, those categories were then grouped into unique pairings for each case. For example, if the toxicology indicated heroin, cocaine, alcohol and fentanyl, then each unique combination would be represented: heroin - cocaine, heroin-alcohol, heroin-fentanyl, cocaine-alcohol, cocaine-fentanyl, and alcohol-fentanyl. For this reason, the combinations displayed are not mutually exclusive, as one fatal overdose may fall into multiple combination categories. Once each case was grouped into unique combinations, those combinations were aggregated to display the frequency of each combination across all OFR cases.

Data represents reviews conducted from January 18, 2018 to November 30, 2023. The cases in this analysis are those reviewed by OFR teams as well as cases abstracted through Dane County's OFR process. The figures presented here do NOT represent a random sample of overdose deaths in Wisconsin.

Last updated: 10/30/2023

This material development was funded by the Cooperative Agreement NU17CE925003-02-02 Centers for Disease Control and Prevention (CDC). Its contents do not necessarily represent the official views of the CDC or the U.S. Department of Health and Human Services. In addition, this project was supported by Grant No. 2018-PM-BX-K101 awarded by the Bureau of Justice Assistance. The Bureau of Justice Assistance is a component of the Department of Justice's Office of Justice Programs, which also includes the Bureau of Justice Statistics, the National Institute of Justice, the Office of Juvenile Justice and Delinquency Prevention, the Office for Victims of Crime, and the SMART Office. Points of view or opinions in this document are those of the author and do not necessarily represent the official position or policies of the U.S. Department of Justice.