## MR-Only or Reversed Workflow Imaging QA Checklist $(9/28/2016\; {\rm EP})$

Patien	t:	RT #:
Date: _		Tumor Site: MD:
Treatn	nent:	New Rescan Boost Scan Orientation: HFS HFP FFS FFP
MR Si	<b>m:</b> (R7	Therapist: MR Technologist:)
Yes	N/A	<u>Item</u>
		Patient setup (Prone breast styrofoam; otherwise flat table overlay)
		Patient orientation ( <b>Brachy:</b> FFS)
		Target volumes centered within MR bore (to extent possible)
		Brain/HeadNeck: Mask and skin are completely dry or mask heated in oven
		Pancreas/Liver/Lung/Esophagus: Headphones molded into alpha cradles/vac-locs
		Abdomen/Chest/Pelvis: Large RF flex coils supported by coil bridges
		Abdomen/Chest/Pelvis: Headphones placed on patient
		Chest: ECG leads placed on patient
		Cholangio/Bile Duct: Nasal cannula placed on patient with oxygen at 2-3 Liters/min
		Cholangio/Bile Duct: Injector loaded with Eovist
		Brachy/Abdomen/Pelvis: 0.5mg glucagon IV at start, midway of exam
		Fiducial markers placed (if not placing isocenter)
		<b>Prostate:</b> Bladder, rectum fill check using scout (call physician to approve)
		Synthetic CT series run as straight axials (switch to sagittal, then switch back to axial)
		High order shim volume (adjust volume) optimized and copied to each series
		B0 map acquired with body coil (not flex coils); magnitude and phase (2 series) in browser
		Additional diagnostic sequences added for MR Sim with Interpretation
		High bandwidths or Advanced WARP used for metal (hip prostheses, spine hardware)
		<b>Brachy:</b> 3D SPACE images acquired as straight axials
		Abdomen/Chest: Breath holds at end expiration
		Coverage sufficient:
		Check EPIC for specific instructions
		• <b>Brachy 3D SPACE:</b> 2cm superior of uterus (page MD)
		Brachy 2D T2s: Cover cervix and uterus
		Prostate: Full bladder coverage
		Sarcoma: 5cm superior/inferior of markers for post-op patients    126
		Images screened for artifacts. Did fat-sat, Dixon separation work? (re-run if necessary)
		Spine/Sarcoma: Ctrl+3,4 to align upper+lower axial groups; merge as 1 series in 3D viewer
		3D distortion correction applied to all images
		3D distortion-corrected images (images with _DIS3D suffix) sent to MIM_Clinical
		Non-distortion corrected images (images with _ND or _DIS2D suffix) deleted from PACS
		Setup reference defined in MIM: BB Iso
	1 1 1	Advanced Imaging: Raw k-space data transferred with Yarra client (Ctrl+Esc → Transfer Raw Data)